

Miso and its biological effects  
Hiromitsu WATANABE, Ph D  
Research Institute for Radiation Biology and Medicine.  
Hiroshima University.  
1-2-3 Kasumi, Kasumi 1-2-3, Minami-ku, Hiroshima, Japan 734-8553

Phone: 81-82-257-1550, Fax 81-82-257-5893

E-mail [tonko@hiroshima-u.ac.jp](mailto:tonko@hiroshima-u.ac.jp)

## **Introduction**

A traditional ingredient of the Japanese diet, miso (fermented soy bean paste), is fermented from soybeans, rice, wheat, or oats. It contains vitamins, microorganisms, salts, minerals, plant proteins, carbohydrates, and fat. Miso also contains saponin inhibiting lipids peroxide, trypsin inhibitor, isoflavon, lecithin, colin, prostaglandin E and others<sup>1)</sup>. It is used on a daily basis as a flavor in soup and solid food in Japan and other parts of Asia and remains an essential ingredient for Japanese-style cooking. Even though miso has no equal in the West, Western cooks familiar with miso prize it for its almost unlimited versatility. It can be used like bouillon, as a rich meat stock in soups and stews<sup>2)</sup>. It is considered as a food with health-promoting benefits, such as effectiveness in relieving fatigue, regulation of the intestinal function, digestive supplement, protection against gastric ulcer, decrease of cholesterol, decrease of blood pressure, whitening ability, prevention of diseases associated with adult lifestyle habits, apoplexia cerebri, accumulation of brain metabolism, protection of aging, healing radiation damage and prevention of cancers for biological effects.

In this review, the effects of miso for radiation prevention, cancer prevention and prevention of hypertension were described with a focus on epidemiological and experimental tests.. .

### **1. Radiation protection by miso**

In August 9<sup>th</sup>, 1945, the 2<sup>nd</sup> atomic bomb was dropped on Nagasaki. At the time, physician Tatuichirou Akizuki, worked with 20 employees, caring for 70 tuberculoses patients in "Uragami Daiichi Hospital" (St Francis Hospital) located about 1.4km away from the hypocenter. However, these people, including Dr. Akizuki, escaped from death caused by acute radiation damage. Dr. Akizuki conjectured that the reason there was no nuclear bomb disease was that these people had consumed cups of wakame miso soup (miso soup with garnish of wakame seaweed)<sup>3)</sup> everyday. Later, his assumption was published in the English language for the information of the Western population. On April 26, 1986, after the accident at Chernobyl, Russia, many Europeans consumed miso soup to prevent radiation diseases. In effect, Dr. Akizuki was the first person in Japan to promote the discovery of the preventive characteristics of miso against radiation damage and also its contribution to a healthier life among the modern generation.

Based on this case of Dr. Akizuki, our experimental studies were started to research the protective effect of miso against radiation damage.<sup>5-7)</sup>

A week before irradiation, five-week-old male B6C3F1 mice were fed a commercial diet of MF (common rodent chow, Oriental Yeast, Tokyo) and MF with 10 % dried red rice miso obtained from Miso Central Institute, Tokyo. The mice for 6 weeks were irradiated with 7 – 12Gy (dose rate 4 Gy/min) for small intestinal crypt survival and were not anesthetized during the irradiation. An histological autopsy was performed on the animals for observation of crypt survival 3.5 days after irradiation.

No radiation protection was shown on 2% of NaCl or MF diet (common rodent diet) groups but the numbers of surviving crypts were significantly increased on miso group. No protective effects were evident when miso was given immediately, 1 or 2 days after irradiation. We concluded that fermented substances might protect not only against gastrointestinal damage but also bone marrow death by radiation. Therefore, it was shown that the protection effectiveness required minimum concentration of the preventive substance(s) at certain level in blood by long term intake of miso<sup>4)</sup>.

An investigation was then conducted to prove whether the protection effect was caused by soybean alone or fermentation periods of miso. One week before irradiation, the mice were given a diet supplemented with 10% of miso at different stages of fermentation such as 3 to 4 days, 120 days and 180 days fermented. After 7 Gy, the irradiation number of surviving crypt was significantly greater for the 120 days fermented miso and 3 to 4 days fermented miso as compared to the MF group. With 8 Gy, surviving crypts in the 3 to 4 days and 180 days fermented miso groups were also significantly increased as compared with the MF values. Crypt survival was significantly different in the 180 days fermented miso group ( $p < 0.01$ ) as compared with the 120 days fermented miso and MF groups (Table 1). The lengths of surviving crypts with 7 and 8 Gy irradiation were increased as compared with those at 0 Gy. Crypt survival was significantly different in the 180 days fermented miso as compared with 3-4 days fermented, 120 days fermented miso and MF groups<sup>6-7)</sup>.

By using miso produced by Miso Central Institute (Tokyo) or Hiroshima Prefectural Food Technology Research Center, crypt survivals were studied at the

dose of 12Gy X-irradiation. There were no significantly different numbers of surviving crypts on both 180 days fermented miso groups. In different types of miso provided from different areas of each institutes, longer fermented miso increased the number of survival crypts and the period of 180 days fermentation is considered more significant to radiation protection. The mechanism of radioprotective effect of miso is considered to be closely related to the substance produced during fermentation stages<sup>8)</sup>. Thus, the radiation protection effects of miso might be attributed to the fermentation process.

The survival rate of the animals at the dose rate of 4 Gy/min did show that there was no difference in groups of mice irradiated at the dose rate of 2 Gy/min. With 15Gy, animals started to die on the 5<sup>th</sup> day. On the 7<sup>th</sup> day, all animals were dead and no significant difference was observed. But with 8 Gy at 10<sup>th</sup> day, MF group started to die on the 5<sup>th</sup> day. At 11<sup>th</sup> day, The 120 days group started to die on the 5<sup>th</sup> day and the 180 days group on the 13<sup>th</sup> day. Using the Cox model, results proved that the 3-4 days ( $p < 0.048$ ), 120 days ( $p < 0.026$ ), and 180 days ( $p < 0.011$ ) groups that were fed the fermented miso had a significantly longer survival period than the MF group. Therefore, at high dose irradiation, the death of the animals was inevitable; but at low dose, after 180 days, the fermented miso has an effect of radiation protection<sup>6)</sup>. We can conclude that the fermented substances protected not only against gastrointestinal but also bone marrow death by radiation. Houchen et al<sup>9)</sup> reported that expression of FGF-2 is induced with radiation injury and recombinant human FGF-2 markedly enhanced crypt survival. Takahama et al<sup>10)</sup> also reported that a replication-deficient adenovirus containing the HST-1 gene acts as a potent protector against lethal irradiation associated with injury to the intestinal tract as well as myelosuppression in the bone marrow and spleen. Ferrel et al<sup>11)</sup> have reported that recombinant human keratinocyte growth factors can protect mice from chemotherapy- and radiation-induced gastrointestinal injury and death but not against whole-body radiation, at least in terms of death from intestinal and contaminated marrow. We also found VEGF to have protective Katoh and Watanabe, (unpublished data). Cytokine-like substances in miso may thus play an important role in the prevention and/or the recovery and repopulation of critical tissue elements when given prior to and during radiation exposure. However, to our knowledge there are no reports regarding extraction of cytokines from miso<sup>9-11)</sup>.

Furthermore, we are conducting a search for effective materials. Water soluble

and ether soluble fraction as well as the residue from 10 days and 260 days fermented miso were investigated. Only water soluble fraction of 10 days miso has no effect on crypt survival but the rest of the fractions of 10 days and 260 days did have an effect on crypt survival. As for mortality, only water soluble fraction of 260 days miso increased survival rate but the rest of each fraction of 10 days and 260 days had no difference compared with control (MF). This suggests that the effective substances in this fraction might be synthesized or the preventative substance in 10 days miso was produced during 260 days fermentation time<sup>8)</sup>.

On the other hands, aglycon type isoflavon and melanoidin were produced during fermentation of miso, so radiation protection of those substances was studied. Aglycon type isoflavon did not have any radiation protective effect. (Watanabe et al. unpublished data) but chemically synthesized melanoidine and the melanoidine-like substance extracted from 7-month fermented miso did affect radiation protection. Therefore, melanoidine like substances in 180 days fermented miso might be considered to have radioprotective effect<sup>12)</sup>. Further study is needed to discover the substances responsible for increased crypt survival, crypt lengths and prolongation of average time to death with miso, their mechanisms and any associated changes in bacterial flora in the gastrointestinal tract.

## **2. Prevention of gastric tumor**

### **2.1 Effects of salt in miso**

Hirayama reported that consuming miso soup everyday reduced the occurrence of stomach cancer<sup>13)</sup>. On the other hand, as miso contains 10- 12 % salt content, it has been considered that high consumption of NaCl greatly increased the occurrence of stomach cancer<sup>14)</sup>. This chapter discusses the difference between salt in miso and salt itself.

Male CD:Crj rats were fed *N*-methyl-*N'*-nitro-*N*-nitrosoguanidine (MNNG) in their drinking water for four months. The rats were simultaneously maintained on a diet supplemented with 10 % dry red miso, 5 % red dry miso, 2.2% NaCl since 10% dry miso contains the equivalent amounts of 2.2% NaCl or 1.1% NaCl and 5 % miso contains 1.1% NaCl or control MF feed. As control, MF in stead of MNNG, tap water is given during diet<sup>15,16)</sup>. Generally, the incidence of gastric tumors of NaCl groups was higher than miso groups. especially, the

incidence of gastric tumors of the MNNG+2.2% NaCl. The size of gastric tumors in the MNNG+2.2% NaCl and MNNG+1.1% NaCl groups and adenocarcinomas in MNNG+1.1% NaCl groups were significantly increased as compared with MNNG+MF. (Table 2)

Furthermore, the same experiment was conducted on 50% salt-reduced miso but the incidence of gastric tumor in the reduced salt content miso (1% of NaCl content) was the same as the regular miso containing 2% NaCl<sup>17)</sup>. This study indicated an increased preventative effect on gastric tumor by miso than by salt diet. Asahara et al.<sup>19)</sup> and Rajendran et al.<sup>20)</sup> reported that by Umu test to indicate mutagenicity, the strains of yeast, lactic acid bacteria and other molds in miso can remove or detoxicate carcinogens as trp-p-2 from burnings of foods. Yanagihara et al.<sup>21)</sup> reported that by using digestive tract cells, a group of isoflavone-like substance in miso such as biochanin A and genistaine prevented the growth of gastric tumor cells and caused the destruction of the cells by apoptosis.

These reports suggested that substances contained in miso destroyed or degenerated carcinogens to decrease the carcinogenicity of MNNG.

## **2.2 Effects on different fermentation stages of miso**

To determine whether soybean themselves or the fermentation process produced the preventive agents, a study was conducted to examine the effects of miso at various fermentation-stages on the initiation phase of MNNG-induction of rat glandular stomach tumors.

Male CD:Crj(SD) rats were maintained on MF diet supplemented with 10 % of dried miso prepared from three different fermentation stages such as 3 to 4 days, 120 days and 180 days with 1% NaCl (Wako, special grade) feed as control. During administration of carcinogen, the animals were fed with the above diets.<sup>22)</sup> (Table 3) There was no significant difference among groups in total number of tumors, number of tumors per rat, incidence of gastric tumors, duodenum tumors and other tumors. For gastric tumors, the size of the tumors in 180 days miso group decreased significantly compared with control (P<0.01). Size of tumors in 3-4 days fermentation miso group were significantly increased as compared with 120 days fermented miso (P<0.05) and 180 days fermented miso (P<0.01). There is a high possibility that during fermentation period of

miso, the substances to inhibit growth of tumors might be produced.

During fermentation of miso, two reactions occur: enzymatic reaction by koji (rice starter with *Aspergillus oryzae*) and fermentation reaction by microorganisms as yeast, lactic acid bacteria. By enzymatic reaction, soybean and rice were decomposed and from hydrolysis of protein, saccharification of starch and decomposition of oil, low molecular peptide, amino acids, reduced sugars, fatty acids and glycerol were produced. A part of these nutrients were consumed by salt-tolerance microorganism, as a by-product of metabolism of fermentation. Alcohol, organic acids and esters, flavor components of miso, were synthesized. In addition, maillard material was produced by non-enzymatic reaction and the material is a factor of color in miso<sup>2)</sup>. Fujinami et al. reported that content of glutamine and formol nitrogen were increased during miso fermentation<sup>22)</sup> and Higashi et al. reported the production of amylase and acid protease during the fermentation period<sup>23)</sup>. These compounds might be produced from fermentation as well as the soy bean.. However, analysis of these compounds is for future investigation.

### **3. Prevention of aberrant crypt foci and of large intestinal cancer**

The recent change of dietary habits from traditional Japanese to Western style diets has been paralleled by an increase in colon and breast cancer in Japan. The change is more responsible for the increase in cancer than genetics.<sup>24)</sup> It was reported that an epidemiologic study showed cancer of the large intestine was suppressed by soy bean<sup>25)</sup>. Many articles had been published concerning physiological property of soy bean inhibiting large intestinal cancer or aberrant crypt foci (ACF) in the aspects of vegetable fiber,<sup>26)</sup> isoflavon,<sup>27-29)</sup> saponin<sup>30)</sup> and high molecular weight fraction of water soluble material<sup>31,32)</sup> of soybean. There also are articles to describe the non-effective aspect of soy bean on cancers<sup>28,29)</sup>.

#### **3-1. Effect of miso, salt, kojic acid and biochanin A on ACF**

The study was designed to investigate the effects of fermented miso, pure NaCl, crude salts, koji acid and biochanin A in the diet on the induction of aberrant crypt foci (ACF) by azoxymethane (AOM), a carcinogen for large intestinal cancer<sup>33-35)</sup>. (Table 3)

Six-week old male F344/DuCrj rats were fed on each test diet.. After one week, AOM (Sigma) was injected by 15 mg/ kg/body once per week for 3 weeks and the rats continued to be fed different diets for 2 weeks. All rats were fed a commercial diet MF alone or with 5%, 10%, 20% of dried red miso, 2.2 % and 4.4% of NaCl (Wako Pure Chemical) which are equivalent NaCl content of 10 % and 20% miso respectively, 4.4% of sea salt (Ryoen, Tokyo), coarse salt (Diasalt, Tokyo), rock salt (Tomen, Tokyo), table salt (JT, Tokyo) and 1.5 % or 3.0% of kojic acid (Nagase Seikagaku), 5 ppm of biochanin A (Sigma). After the animals were euthanized, the colon was removed and fixed in 10% buffered formalin and stained with 0.5% methylene blue. By viewing the stained colons with a light microscope at magnification of 20-30 times, the numbers of ACF, total number of aberrant crypt (AC) per colon and of AC per focus were counted.

All rats treated with AOM showed 100% incidence of ACF. The number of ACF per animal was significantly decreased except in the groups of 4.4 % pure NaCl, 2.2 % pure NaCl group, 5 % miso and biochanine A. The number of total AC significantly decreased except in the group of 2.2% of pure NaCl, 3% kojic acid and biochanine A. AC per focus increased significantly in the 10%, 20% miso group.

ACF decreased as concentration of miso increased but not by the pure NaCl. ACF decreased in the group of crude salts and 3.0% of kojic acid but not in the group of 1.5 % kojic acid and biochanine A. Kojic acid exists in miso during fermentation period but does not in the final product. (Mohri & Fujinami, personal communication). Therefore, the suppressing effect on ACF of miso did not appear to be kojic acid nor similar substance.

This study showed pure NaCl did not have a suppressing effect on ACF but crude salt did. Some of the groups were measured by BrdU labeling index or germinal regions but there was no correlation between ACF and AC induction. Impurities of purified salt used in this study were less than 0.01% but crude salt contained calcium and minerals and these substances were reported to suppress cancer in the large intestine and ACF induction<sup>36-41)</sup> while also preventing DNA synthesis<sup>42,43)</sup>. It is also reported that seren<sup>46)</sup>, copper and iron<sup>47)</sup> had the same effect of suppression. Therefore, suppression effect of ACF induced by AOM was considered to be caused by the minerals in miso such as calcium and

magnesium. This fact suggests that large intake of minerals in sea salt used in Japanese regular food, is a factor in lowering the occurrence of large intestinal cancer among Japanese people.

### **3-2 Effects of long term fermented miso on ACF**

The study was designed to investigate the effect of different fermentation times of miso on the ACF induced by AOM<sup>46)</sup>.

All rats treated with AOM developed ACF. The number of ACF per animals were,  $87.8 \pm 28.9$  in group of MF,  $84.9 \pm 52.1$  in 2.2% NaCl,  $85.5 \pm 19.3$  in short-term (3-4 days) fermented miso,  $83.6 \pm 20.8$  in middle-term (120 days) fermented miso,  $65.1 \pm 18.4$  long-term (180 days) fermented miso. The number of ACF in long-term miso decreased significantly compared with MF and short term, and middle term groups. Total AC number in middle-term miso was largest and increased significantly compared with 180 days miso. In this study, long term fermented miso was the most effective suppresser of ACF induced by AOM. The effect was considered to be caused by components of soy bean, minerals, and other substances produced during fermentation of miso as previously described.

### **3-3. Effect on large intestinal adenocarcinomas**

Five-week-old male F344/DuCrj rats were maintained on MF diet supplemented with 10 % of each dried miso prepared from two different fermentation stages such as short term (3 to 4 days) and long term (180 days). Twenty-four weeks after administration of carcinogen AOM, the animals were euthanized and the colons were investigated. There was no statistical difference among the groups in averaged number of ACF per colon, total number of AC and number of AC per ACF<sup>47-48)</sup>. (Table 4)

Each animal that died on 103<sup>rd</sup>, 149<sup>th</sup> and 163<sup>rd</sup> day after the first AOM injection in miso groups had signet cell carcinoma in the duodenum. By pathological observation, signet ring cell carcinomas in the large intestine were not observed in the group of 180 days fermented miso. The size of well-differentiated types of adenocarcinomas and total adenocarcinomas (well-differentiated type and signet ring cell) in the 180 days and 3-4 days fermented miso groups was significantly smaller in MF+AOM group. The

number of crypts cell, PCNA-positive cells, the highest position and width of the germinal region of positive index, decreased significantly in 180 days miso compared with MF+AOM values. In the present study, dietary administration of 180 days fermented miso inhibits the development of AOM-induced ACF in the rat colon. But 10 % miso did not have the suppressive effect on ACF induced by DMH<sup>50)</sup>.

### **3-4. Effects of soybean products on colon carcinogenesis**

Epidemiological study showed that intake of miso soup decreased large intestinal cancer<sup>50), 51)</sup>. Kono et al reported that taking two cups of miso soup reduced the amount of S type colon cancer.<sup>52)</sup> By taking tofu and soybean, Watanabe et al reported a decrease in rectal tumors<sup>53)</sup> and Poole et al reported a decrease in large intestinal tumors<sup>54)</sup>

caused by soybean products and miso. Hu et al reported a decrease in colon tumors of men in China<sup>55)</sup>. LeMarchand et al reported, intake of soybean or soybean products in Hawaii reduced occurrence of large intestinal cancer in women but not in men<sup>56)</sup>. Tuyns et al reported soybean to be clearly a preventive agent against colon rectal cancers on the basis of a case control study in Belgium<sup>57)</sup>.

In the study by Davies, F344 rats were fed on soybean with high content isoflavon and on a diet supplemented with high fat and with less calcium one week before and 31 weeks after AOM administration. The soybean group showed no decrease in the occurrence of large intestinal tumors<sup>58)</sup>. Min mice were fed a Western type of diet (high fat and less vegetable fibers and calcium) with higher amount of isoflavon added, but the study did not show any preventive effect on large intestinal cancer<sup>59)</sup>. Rao<sup>60)</sup> reported the increase in non-invasive tumor numbers and adenocarcinomas based on the study of AOM-treated rats fed by casein-based feed with genistaine. McIntosh<sup>61)</sup> reported a greater increase in the number of tumors with soy protein than casein. Gee<sup>62)</sup>, using a DMH model, reported no suppression of ACF in large intestinal tumors by genistein. I. Hahhak et al<sup>63)</sup> and Thiagarajan<sup>64)</sup> observed a suppressive effect of soy or soy products against ACF induction. Pereira et al<sup>65)</sup> also found purified genistein to inhibit induction of ACF. Since isoflavon has anti-estrogen activity, it is considered to be a most effective substance against breast cancer<sup>66-68)</sup>. Gotoh et al demonstrated that administration of

biochanin A, a genistein precursor, inhibited the development of mammary tumors but we reported biochanin A did not reduce colonic ACF development in rats treated with AOM<sup>69)</sup>. It is considered that isoflavon has a preventive effect on cancer of breasts and internal organs related to hormone. There is no clarification of the epidemiological relationship between large intestinal cancer and isoflavon.

In the present study, long-term as 180 days fermentation miso significantly decreased in the PCNA positive index. It is hypothesized that calcium is a regulator of cell proliferation in the colon. Dietary intake of calcium inhibits experimental carcinogenesis<sup>70)</sup>. Therefore, during fermentation of miso, the substance to suppress cell proliferation is generated and it might work as a factor in preventing colonic cancer

#### **4. Prevention of lung cancer**

In Japanese, lung cancer has exceeded stomach cancer and become the highest incidence of cancer. Since it became a serious problem, it is an urgent matter to develop a preventive method of lung cancer. Epidemiological studies in Okinawa<sup>71)</sup> and Singapore<sup>72)</sup> have indicated that fermented soy-bean products might have inhibitory effects on lung cancer. Shiraki<sup>73,74)</sup> investigated such effects of fermented miso by using short-term fermented as 3 -4 days, and long-term as 180 days miso.

Six-week old male Slc:Wister rats were treated di-isopropanol-nitrosamine (BHP) in drinking water at concentration of 200 mg/liter for 10 weeks. All rats were fed by commercial diet MF (Oriental Yeast Co., Ltd. Tokyo) alone or with added miso. Short-term dry red miso, 3-4 days, and long-term, 80 days, fermentation (Miso Central Institute, Tokyo) were supplemented into MF at 10%. The diets were supplied with normal tap water *ad libitum* for 12 weeks after treatment of BHP and the rats were euthanized 22 weeks after the treatment. Macroscopically, all animals had many nodules in the lungs. The number of nodules in the 180 days fermented miso group was significantly lower as compared to other groups but the size of tumors did not show significant difference among groups. All animals had hyperplasia, adenomas and adenocarcinomas. Papillomas or squamous cell carcinomas were also evident in

13%, 27% and 29% of the animals in the BHP, BHP+short, BHP+long but rate of incidence had no significant different. numbers of adenocarcinoma in the BHP+long and PCNA. Strongly positive tumors in the miso group were significantly decreased as compared to the BHP alone group. Size of tumors decreased significantly in BHP+long group. It was obvious that the long-term fermented miso reduced the numbers of lung tumors. (Table 6)

Koo et al<sup>75)</sup> reported a significant inverse association between tofu/soy intake and lung cancer in non-smokers in Hong Kong after adjustment for age, numbers of live birth and schooling. Swanson et al<sup>76)</sup> reported a dose-dependent inverse relationship with tofu and risk of lung cancer in Yunnan Province, China. On the other hand, Ozawa et al reported that among females, a high intake of miso-soup significantly and almost dose-dependently increased the risk<sup>77)</sup> However, consumption of miso-soup seems to be associated with Japanese-style diet and increased salt intake. But, the significance of these finding is unclear.

### **5. Prevention of hepatic tumor**

When C3H male mice were bred for 1 year under normal condition, 89% of the mice had natural occurring liver cancer. When B6C3F1 mice were irradiated by <sup>252</sup>Cf neutron, which is the same neutron of atomic bomb, 2Gy. Thirteen months later, 62% of the male mice and 29 % of female had liver cancer. However, these mice were fed on diets containing 10% miso (Miso Central Institute, Tokyo), the frequency of cancer of C3H, male B6C3F1 and female B6C3F1 was reduced to 32%, 13%, 13% respectively<sup>78)</sup>.

Ogundigie et al demonstrated the same effect of miso. Strong carcinogen for hepatocytes, diethyl-nitrosamine (DEN), was injected to 15 days-old B6C3F1 mice and at 4 weeks-old, the mice were irradiated at 2Gy of <sup>252</sup>Cf neutron and at 40 weeks-old, the animals were euthanized for the investigation. Various kinds of preventive materials including miso (Miso Central Institute, Tokyo), were applied to find out occurrence of hepatic tumors. All animals had hepatomas and control groups without any preventive material had 46 tumors/mouse. But in case of miso, number of tumors reduced to be 32.5/mouse significantly. When 10 ppm and 20 ppm of biochanin A were applied, the numbers of 10 ppm were 40.1/ mouse but the number of 20 ppm reduced significantly to be 32.5/mouse<sup>79)</sup>.

Obviously, both results indicated that the administration of miso inhibited the development of hepatomas induced by natural or <sup>252</sup>Cf neutron or carcinogen in

experimental animals.

Kurosawa et al<sup>80)</sup> reported that miso-soup among women without history of liver diseases showed a significant inverse association with hepatocellular carcinoma mortality. And also Sharp et al<sup>81)</sup> suggested that consumption of soy foods especially miso-soup, is associated with reduced risk of hepatocellular carcinoma.

## 6. Prevention of breast cancer

Yamamoto et al<sup>82)</sup> reported that taking a cup of miso three times a day had reduced the occurrence of breast cancer but tofu, natto, soybean and fried bean curd did not have such effect and women in menopause had less breast cancer when a large amount of isoflavon was taken.

Baggott et al reported that when dimethylbenz[a]-anthracene (DMBA) was administered to Sprague-Dawley (SD) rats, 4.1 per rats of mammary tumors occurred but when the animals were fed on diets containing 25% miso, the numbers of tumors were reduced to be 2.95 per rat and the time until occurrence of tumor with miso became longer than with only carcinogen<sup>83)</sup>.

Gotoh et al<sup>69,84)</sup> investigated the induction of mammary tumors by administration of soybean, miso, and biochanin A in 40mg/kg of *N*-nitroso-*N*-methylurea (MNU) treated SD rats. The incidence time of the tumors was slower in miso and biochanin groups than MNU itself. The number of tumors was 2.2 in MNU but in miso group, it significantly reduced to be 1.2 and also in 10 % soybean and biochanin A groups. They also investigated the incidence of tumors by miso and tamoxifen, a medicine used for breast cancer treatment. The number of tumors per rat induced by MNU only was 4.5 but it was 2.4 in miso groups and it was 1.4 in tamoxifen groups and the number was significantly reduced to be 0.2 when miso and tamoxifen were fed together. This investigation clearly indicated that administration of miso in the diet can reduce the occurrence of mammary tumors as well as tamoxifen at the beginning of carcinogen treatment and it can also suppress the multiplicity of tumors induced by carcinogen.

## 7. Protection against hypertension.

It was often considered that salt in miso induced stomach cancer<sup>14)</sup> as well as the

increase in blood pressure<sup>85-91)</sup> and even excess amount of intake of miso led to lifestyle-related disease. As described before, it is unlikely salt itself or salt in miso did not cause the increase of gastric tumor. Afterwards, the effects of salt in miso on blood pressure were studied<sup>92)</sup>.

Six-week-old salt-sensitive Dahl rats or similar SD male or female rats (original strain of salt-sensitive Dahl rat) were maintained on commercial diet MF alone or supplemented with 10 % dry red miso (180 days fermented miso, Miso Central Institute, Tokyo, Japan) or 2.3% salt because 10% dry miso containing 2.3 % NaCl or 1.9 % salt or 0.3 % salt for 12 weeks. Systolic blood pressure in the Dahl male rats on 2.3% NaCl was significantly increased for 8 and 12 weeks but not in SD rats. However, the blood pressure on diet with miso or commercial control diet (MF) was not increased. Even though miso contains 2.3% NaCl, the blood pressure was stable as on commercial diet containing 0.3% salt. (Table 7)

Systolic blood pressure of Dahl female rats was increased on salt diet but not increased on miso diet. The blood pressure of female rats was not increased on 1.9% salt diet but that of male was increased. Male rats are more sensitive to salt than female rats similar to human beings. (Table 8)

As a result, blood pressure of salt-sensitive rats was not increased on miso diet that included 2.3% NaCl. This indicated that these rats are regarded as salt-sensitive as human beings. It was concluded that the miso diet did not increase blood pressure in such people. Salt in miso worked in different manner from salt itself for gastric tumors as well as blood pressure. Considering the result of blood pressure between salt in miso and salt itself, it is better to refrain from taking a lot of salt. Rather it is recommended to take miso soup more often to prevent the occurrence of lifestyle-related disease.

In epidemiological data, He et al demonstrated that dietary soybean protein reduced blood pressure in a randomized, double-blind, controlled trial in Chinese adults<sup>86,87)</sup>. Several small clinical trials have reported inconsistent findings regarding the effect of soybean protein on blood pressure<sup>91,94,95)</sup>. Washburn and colleagues<sup>91)</sup> compared the effect of 20 g of soybean protein containing 34 mg of phytoestrogens given either in 1 dose or in 2 doses with that of 20 g of complex carbohydrates on cardiovascular disease risk factors and menopausal symptoms among 51 women in a randomized, controlled trial.

They observed a significant reduction in diastolic blood pressure in the twice-daily soybean protein diet compared with the carbohydrate control diet. Burke and colleagues<sup>94)</sup> examined the effects of soybean protein on 24-hour ambulatory blood pressure among 41 treated hypertensive patients in a randomized, controlled trial. He also documented that blood pressure reduction associated with soybean dietary protein was greater than that of the carbohydrate control<sup>92)</sup>. Conventionally, increased intake of complex carbohydrate has been recommended as a replacement for saturated fat intake to reduce cardiovascular risk<sup>96)</sup>. Soy-based diet attenuated the development of hypertension both in female and male SHR rats<sup>97)</sup>. These findings suggested that increased intake of soybean protein may play an important role in prevention and treatment of hypertension.

Nakamura et al<sup>97)</sup> conducted a double-blind, randomized placebo-controlled study to evaluate the effects of 6-week diet containing approximately 25% to 20% low-sodium soy sauce and miso (an approximately 10% reduction of total dietary salt intake) in Japanese on blood pressure. The changes in blood pressure observed in the experiment were not significant. However, in those aged 40 years and older, a 6.4mm Hg net reduction in diastolic blood pressure with no significant change in systolic blood pressure was noted in the low-sodium intake group. Nakamura et al reported that a 6-week trial of low-sodium soy sauce and miso resulted in a reduction in urinary salt of 0.7g/day in the intervention group and the net difference of urinary salt excretion between the intervention group and the controls was 1.4g/day. They concluded that an approximately 10 % reduction of dietary salt intake affected diastolic blood pressure in middle-aged or older people. When one consumes 13g of dietary salt daily, the estimated salt intake from soy sauce will change from 3.3g to 2.4g and that from miso will change from 1.6 to 1.3 by replacing the seasoning with the low-sodium type. Thus one can achieve 1.2g reduction of daily dietary salt intake. Dietary reduction of NaCl in soy sauce rather than miso may be used to reduced diastolic blood pressure. On the other hand, Kanda et al performed a study of the association of lifestyle parameters with the future risk of hypertension in normotensive subjects. They administered a baseline questionnaire and performed a four-year follow-up of 445 normotensive Japanese from 35 to 89 years of age. In 60 to 69 year old subjects, changes of blood pressure during four years were negatively correlated with boiled rice intake in men and with Japanese tea intake in women<sup>98)</sup>. Multiple logistic regression analysis revealed that miso-soup intake of two or more bowls per day

protected against hypertension during the follow-up ( $p < 0.05$ ). These results indicate that the nature and amounts of food intake is important in the prevention of hypertension in the elderly. It also prevents increased blood pressure when compared with rats fed the equivalent NaCl concentration without miso in men and animals. For protection of hypertension, Fujita recommended<sup>99)</sup> that taking miso soup together with protein or together with potassium rich vegetable as spinach or wakame seaweed prevented the increase in blood pressure even in salt sensitive persons.

### **Conclusion;**

Highest occurrence of cancer used to be stomach cancer in Japan. However, the recent change of dietary habits from traditional Japanese to Western style increased the occurrence of cancer of large intestine, breast cancer and prostate cancer more than stomach cancer. This indicates the importance of dietary habit and traditional dietary habits as miso soup should be re-evaluated not only in Japan but in the world. Since miso has such functions as discussed in previous sections for preventive effect on maintaining your health, further studies are needed to clarify the mechanism and effective materials produced during fermentation in miso. As a whole, miso is considered to be a good food ingredient for health.

Finally, the effective substances caused during fermentation are unclear. Kennedy<sup>100,101)</sup> have declared that soybean-derived Bowman-Birk inhibitor suppresses carcinogenesis. This substance should be one of the candidates for analysis of biological effective compounds in miso. This is for future investigation.

### **References**

- 1) Miso Health Promotion Committee: [www:miso.or.jp/miso-e/index/html](http://www.miso.or.jp/miso-e/index/html)
- 2) Shurtleff W, Aoyagi, A: The book of miso, Autumn Press, 1976
- 3) Akizuki S: Health condition and diet , Kurie press, 1980 (in Japanese)
- 4) Watanabe, T, Takahashi, T, Ishimoto, T, Ito, A: The Effect of miso diet on small intestinal damage in mice irradiated by X-ray. Miso Tech Sci. 39, 29-32, 1991. (in Japanese)
- 5) Watanabe T, Ito A: Radioprotective effects of miso in digestive tract and preventive effect of miso on tumors. Daizu Geppou, November, 11-17,1996.

(in Japanese)

- 6) Ohara M, Lu H, Shiraki K, Ishimura Y, Uesaka T, Katoh O, Watanabe H: Radioprotective effects of miso (fermented soy bean paste) against radiation in B6C3F1 mice: increased small intestinal crypt survival, crypt lengths and prolongation of average time to death. *Hiroshima J Med Sci*, 50, 83-86, 2001.
- 7) Ohara M, Huimei Lu, Hamda H, Nishioka T, Kageyama N, Ishimura Y, Shiraki K, Uesaka T, Katoh O, Watanabe H, Kawano K: Radioprotective effects of long-term fermented miso against irradiation in B6C3F1 mice. *Miso Tech Sci*. 50, 21-27, 2002. (in Japanese)
- 8) Kawano k, Matsuda S, Kashiwabara S, Kashimoto N, Kageyama N, Hashimoto K, Nishioka T, Uesaka T, Katoh O, Watanabe H: Effect of each fractions extracted from miso on the protection against radiation damage and aberrant crypt foci. *Miso Tech Sci*. 51, 429-434, 2003.(in Japanese)
- 9) Houchen CW, George RJ, Strumoski MA, Cohen SM: FGF-2 enhances intestinal stem cell survival and its expression is induced after radiation injury. *Amer J Physiol*, 276, G249-G258, 1999.
- 10) Takahama Y, Ochiya T, Tanooka H, Yamamoto H, Sakamoto H, Nakano H, Terada M: Adenovirus-mediated transfer of HST-1/FGF-4 gene protects mice from lethal irradiation. *Oncogene*, 18, 5943-5947, 1999.
- 11) Farrell CL, Bready JV, Rex KL, Chen JN, DiPalma CR, Whitcomn KL, Yin S, Hill DC, Wiemann B, Starnes CO, Havill AM, Lu Z-N, Aukerman SL, Aukerman L, Pierce GF, Thomason A, Potten CS, Ulich TR, Lacey DL: Keratinocyte growth factor protects mice from chemotherapy and radiation-induced gastrointestinal injury and mortality. *Cancer Res*, 58, 933-939, 1998.
- 12) Watanabe T, Shiono T, Kawamura D, Kawano K, Kajimura J, Kashimoto N, Kamtani K; Radioprotective effects of melanoidine against irradiation in B6C3F1 mice. *J Nagasaki Med Ass* in press. (in Japanese)
- 13) Hirayama T: A large scale cohort study on cancer risks by diet with special reference to the risk reducing effects of green yellow vegetable consumption. In *Diet, Nutrition and Cancer*, eds Hayashi Y, Nagao M, Sugimura T, Takayama S, Tomatis L, Wattenberg LW, Wogan GH, pp.41-53, Japan Sci. Soc. Press and Utrecht Vnu Sci Press, Tokyo, 1986.
- 14) Tsugane S, Sasazuki S, Kobayashi M, Sasaki S, for the JPHC Study Group: Salt and salted food intake and subsequent risk of gastric cancer among middle-aged Japanese men and women. *Br J Cancer*, 90,128-134, 2004.
- 15) Watanabe H, Uesaka T, Kido S, Ishimura Y, Shiraki K, Kuramoto K, Hirata S,

- Shoji S, Katoh O, Fujimoto N: Influence of concomitant Miso or NaCl treatment on induction of gastric tumors by N-methyl-N'-nitro-N-nitrosoguanidine in rats. *Oncol Rep*, 6, 989-993, 1999.
- 16) Watanabe T, Ishimoto T, Okamoto T, Takahashi T, Ito A: Effects of miso diet or sodium chloride diet on gastric tumorigenesis in rats. *Miso Tech Sci.* 40, 324-329, 1992.(in Japanese)
  - 17) Watanabe T, Hamada H, Nishioka T, Uesaka T, Huimei Lu, Ohara M, Shoji S, Katoh O: Reduction of gastric tumors by administration with reduced salt miso diet during N-methyl-N'-nitro-N-nitrosoguanidine (MNNG) treatment in rats. *Miso Tech Sci.* 48, 22-26, 2000. (in Japanese)
  - 18) Asahara N, Zhang XB, Ohta Y: Anti-mutagenic and mutagen-binding activation of mutagenic pyrolyzate by microorganisms isolated from Japanese miso. *J Sci Food Agric*, 58, 395-401, 1992.
  - 19) Rajendran R, Ohta Y: Binding of heterocyclic amines by lactic acid bacteria from miso, a fermented Japanese food. *Can J Microbiol*, 44, 109-115, 1998.
  - 20) Yanagihara K, Ito A, Tuge T, Numoto M: Antiproliferation effects of isoflavones on human cancer cell lines established from the gastrointestinal tract. *Cancer Res*, 53, 5815-5821, 1993.
  - 21) Ohara M, Lu H, Shiraki K, Ishimura Y, Uesaka T, Katoh O, Watanabe H: Inhibition by long-term fermented miso of induction of gastric tumor by N-methyl-N'-nitro-N-nitrosoguanidine in CD (SD) rats. *Oncol Rep*, 9, 613-616, 2002
  - 22) Fujinami H, Mochizuki T, Sagawa I, Mori M: Change of glutamic acid, glutamine and pyroglutamic acid during fermentation of miso at different temperature on the ripening of miso (I) *J.Brewing Society of Japan* 78, 466-474, 1983.
  - 23) Higashi K: Fermentation and brewing (1), 130, 2002 p130. 2002. Korin Press (in Japanese)
  - 24) Tajima K, Hirose K, Nakagawa N, Kuriishi T, Tominaga, S: Urban-rural difference in the trend of colorectal cancer mortality with special reference to the subsites of colon cancer in Japan. *Jpn J Cancer Res*, 76, 717-728, 1985.
  - 25) Tuyns AJ, Kaaks R, Haelterman M: Colorectal cancer and the consumption of foods: a case-control study in Belgium. *Nutr Cancer*, 11, 189-204, 1988.
  - 26) Barnes DS, Clapp NK, Scott DA, Oberst DL, Berry SG: Effects of wheat, rice corn, and soybean bran on 1,2-dimethylhydrazine-induced large bowel tumorigenesis in F344 rats. *Nutr Cancer*, 5, 1-9, 1983.

- 27) Messine M, Bennink M: Soyfoods, isoflavones and risk of colon cancer: a review of the in vitro and in vivo data. *Ballieres Clin Encocrinol Metab*, 12, 707-728, 1998.
- 28) Davies MJ, Bowey EA, Adlercreutz H, Rowland IR, Rumsby PC: Effects of soy or rye supplementation of high-fat diets on colon tumour development in azoxymethane-treated rats. *Carcinogenesis*, 20, 972-931, 1999.
- 29) Gee JM, Noteborn HPJM, Polley ACJ, Johnson IT: Increased induction of aberrant crypt foci by 1,2-dimethylhydrazine in rats fed diets containing purified genistein or genistein-rich soya protein. *Carcinogenesis*, 21, 2255-2259, 2000.
- 30) Koratkar R, Rao AV: Effect of soya bean saponines on azoxymethane-induced preneoplastic lesions in the colon of mice. *Nutr Cancer*, 27, 206-209, 1997.
- 31) Azuma N, Suda H, Iwasaki H, Kanamori R, Iwami K: Soy bean crude refuse alleviates experimental tumorigenesis in rat colon. *Biosci Biotechnol Biochem*, 63, 2256-2258, 1999.
- 32) Azuma N, Machida K, Saeki T, Kanamoto R, Iwami K: Prevention effect of soybean resistant proteins against experimental tumorigenesis in rat colon. *J Nutr Sci Vitaminol*, 46, 23-29, 2000.
- 33) Masaoka Y, Watanabe H, Katoh O, Dohi K: Effects of miso and NaCl on the development of colonic aberrant crypt foci induced by azoxymethane in F344 rats. *Nutr Cancer*, 32, 25-28, 1998.
- 34) Masaoka Y, Katoh O, Watanabe H: Inhibitory effects of crude salts on the induction and development of colonic aberrant crypt foci in F344 rats given azoxymethane. *Nutr Cancer*, 37, 78-81, 2000.
- 35) Masaoka Y, Hamada H, Nishioka T, Huimei Lu, Ohara M, Ishimura Y, Shiraki K, Katoh O, Watanabe H: Inhibition effects of Aberrant crypt foci (ACF) induced by Azoxymethane(AOM) on miso and the related substances in F344 rats. *Miso Tech Sci*. 48, 393-398, 2000.(in Japanese)
- 36) Wargovich MJ, Lynch PM, Leven B: Modulating effects of calcium in animal models of colon carcinogenesis and short-term studies in subjects at increased risk for colon cancer. *Am J Clin Nutr*, 54, 202s-205s, 1991.
- 37) Newmark HL, Lipkin M: Calcium, vitamin D, and colon cancer. *Cancer Res*, 52, 2067s-2070s, 1992.
- 38) Vinas-Salas J, Biendicho-Felis C, Miguelsanz-Garcia S, Perez-Holanda S: Calcium inhibits colon carcinogenesis in an experimental model in the rat. *Eur J Cancer*, 34, 1941-1945, 1998.
- 39) Baron JA, Beach M, Mandel JS, van Stolk RU, Haile RW, Sandler RS,

- Rothstein R, Summers RW, Snover DC, Beck GJ, Bond JH, Greenberg ER: Calcium supplements for the prevention of colorectal adenomas. Calcium Polyp Prevention Study Group. *N Engl J Med*, 340, 101-107, 1999.
- 40) Marcus PM, Newcomb PA: The association of calcium and vitamin D, and colon and rectal cancer in Wisconsin women. *Int J Epidemiol*, 27, 788-793, 1998.
  - 41) Tanaka T, Shinoda T, Yoshimi N, Niwa K, Iwata H, Mori H: Inhibitory effect of magnesium hydroxide on methylazoxymethanol acetate-induced large bowel carcinogenesis in male F344 rats. *Carcinogenesis*, 10, 613-616, 1989.
  - 42) Mori H, Tanaka T, Sugie S, Yoshimi N, Kawamori T, Hirose Y, Ohnishi M: Chemoprevention by naturally occurring and synthetic agents in oral, liver, and large bowel carcinogenesis. *J Cell Biochem (Suppl)*, 27, 35-41, 1997.
  - 43) Li H, Kramer PM, Lubet RA, Steele VE, Kelloff GJ, Pereira MA: Effect of calcium on azoxymethane-induced aberrant crypt foci and cell proliferation in the colon of rats. *Cancer Lett*, 124, 39-46, 1998.
  - 44) Feng Y, Finley JW, Davis CD, Becker WK, Fretland AJ, Hein DW: Dietary selenium reduces the formation of aberrant crypts in rats administered 3,2'-dimethyl-4-aminobiphenyl. *Toxicol Appl Pharmacol*, 157, 36-42, 1999.
  - 45) Davis CD, Feng Y: Dietary copper, manganese and iron affect the formation of aberrant crypts in colon of rats administered 3,2'-dimethyl-4-aminobiphenyl. *J Nutr*, 129, 1060-1067, 1999.
  - 46) Ohara M, Lu H, Shiraki K, Ishimura Y, Uesaka T, Katoh O, Watanabe H: Prevention by long-term fermented miso of induction of colonic aberrant crypt foci by azoxymethane in F344 rats. *Oncol Rep*, 9, 69-73, 2002.
  - 47) Ohuchi Y, Myojin Y, Shimamoto F, Kashimoto N, Kamiya K, Watanabe H: Decrease in size of azoxymethane induced colon carcinoma in F344 rats by 180-day fermented miso. *Oncol Rep*, 14, 1559-1564, 2005.
  - 48) Ohuchi Y, Myoujin Y, Kashimoto N, Kajimura J, Kamiya K, Watanabe H: Prevention by 180-day fermented miso of colon carcinoma induced by Azoxymethane in F344 rat. *Miso Tech Sci*. 54, 139-148, 2006.(in Japanese)
  - 49) Ishimoto T, Watanabe T, Okamoto T, Matsuda M, Ito A: Effect of miso diet on large intestinal tumorigenesis in mic. *Miso Tech Sci*. 40, 447-451, 1992.(in Japanese)
  - 50) Nishi M, Yoshida K, Hirata K, Miyake H: Eating habits and colorectal cancer. *Oncol Rep*, 4, 995-998, 1997.
  - 51) Hoshiyama Y, Sekine T, Sasabe T: A case-control study of colorectal cancer and its relation to diet, cigarettes, and alcohol consumption in Saitama Prefecture, Japan. *Tohoku J Exp Med*, 171, 153-165, 1993.

- 52) Kono S, Shinchi K, Ikeda N, Yanai F, Imanishi K: Physical activity, dietary habits and adenomatous polyps of the sigmoid colon: a study of self-defense officials in Japan. *J Clin Epidemiol*, 44, 1255-1261, 1991.
- 53) Watanabe T, Tada M, Kawamoto K, Uozumi N, Kajiwara Y, Hayashi K, Yamaguchi K, Murakami K, Misaki F, Akasaka T, Kawai K; A case control study of cancer of the rectum and colon. . *Nisshoushi* 81, 185-193, 1984.(in Japanese)
- 54) Poole C: A case-control study of diet and colon cancer. Boston MA, Harvard School of Public Health, 1989.
- 55) Hu JF, Liu YY, Yu YK, Zhao TZ, Liu SD, Wang QQ: Diet and cancer of the colon and rectum: a case-control study in China. *Int J Epidemiol* 20: 362-367, 1991.
- 56) Le Marchand L, Hankin JH, Wilkens LR, Kolonel LN, Englyst HN, Lyu LC: Dietary fiber and colorectal cancer risk. *Epidemiology*, 8, 658-665, 1997.
- 57) Tuyns AJ, Kaaks R, Haelterman M: Colorectal cancer and the consumption of foods: a case-control study in Belgium. *Nutr Cancer*, 11, 189-204, 1988.
- 58) Davies MJ, Bowey EA, Adlercreutz H, Rowland IR, Rumsby PC: Effects of soy or rye supplementation of high-fat diets on colon tumour development in azoxymethane-treated rats. *Carcinogenesis*, 20, 927-931, 1999.
- 59) Sorensen, IK, Kristiansen E, Mortensen A, Nicolaisen GM, Wijnands JA, van Kranen HJ, van Kreijl CF: The effect of soy isoflavones on the development of intestinal neoplasia in ApcMin mouse. *Cancer Lett*, 130, 217-225, 1998.
- 60) Rao CV, Wang CX, Simi B, Lubet R, Kelloff G, Steele V, Reddy BS.: Enhancement of experimental colon cancer by genistein. *Cancer Res*, 57, 3717-3722, 1997.
- 61) McIntosh GH, Regester GO, Le Leu RK, Royle PJ, Smithers GW: Dairy proteins protect against dimethylhydrazine-induced intestinal cancers in rats. *J Nutr*, 125, 809-816, 1995.
- 62) Gee JM, Noteborn HP, Polley AC, Johnson IT: Increased induction of aberrant crypt foci by 1,2-dimethylhydrazine in rats fed diets containing purified genistein or genistein-rich soya protein. *Carcinogenesis*, 21, 2255-2259, 2000.
- 63) Hakkak R, Korourian S, Ronis MJ, Johnston JM, Badger TM: Soy protein isolate consumption protects against azoxymethane-induced colon tumors in male rats. *Cancer Lett*, 166, 27-32, 2001.
- 64) Thiagarajan DG, Bennink MR, Bourquin LD, Kavas FA: Prevention of precancerous colonic lesions in rats by soy flakes, soy flour, genistein, and calcium. *Am J Clin Nutr*, 68 (Suppl), 1394S-1399S, 1998.

- 65) Pereira MA, Barnes LH, Rassman VL, Kelloff GV, Steele VE: Use of azoxymethane-induced foci of aberrant crypts in rat colon to identify potential cancer chemopreventive agents. *Carcinogenesis*, 15, 1049-1054, 1994.
- 66) Martin PM, Horwitz KB, Ryan DS, McGuire WL: Phytoestrogen interaction with estrogen receptors in human breast cancer. *Endocrinol*, 103, 1860-1867, 1978.
- 67) Lamartiniere CA: Protection against breast cancer with genistein: a component of soy. *Am J Clin Nutr*, 71(Suppl), 1705S-1707S, 2000.
- 68) Sarkar FH, Li Y: Mechanisms of cancer chemoprevention by soy isoflavone genistein. *Cancer Metastasis Rev*, 21, 265-280, 2002.
- 69) Gotoh T, Yamada K, Yin H, Ito A, Kataoka T, Dohi K: Chemoprevention of *N*-nitroso-*N*-methylurea-induced rat mammary carcinogenesis by soy foods or biochanin A. *Jpn J Cancer Res*, 89, 137-142, 1998.
- 70) Appleton GVN, Davies PW, Bristol JB, Williamson RCN: Inhibition of intestinal carcinogenesis by dietary supplementation with calcium. *Br J Surg*, 74, 523-525, 1987.
- 71) Wakai K, Ohno Y, Genka K, Ohmine K, Kawamura T, Tamakoshi A, Lin Y, Nakayama T, Aoki K, Fukuma S: Risk modification in lung cancer by a dietary intake of preserved foods and soyfoods: findings from a case-control study in Okinawa, Japan. *Lung Cancer*, 25, 147-159, 1999.
- 72) Seow A, Poh WT, Teh M, Eng P, Wang YT, Tan WC, Chia KS, Yu MC, Lee HP: Diet, reproductive factors and lung cancer risk among Chinese women in Singapore: evidence for a protective effect of soy in nonsmokers. *Int J Cancer*, 97, 365-371, 2002.
- 73) Shiraki K, Une K, Yano R, Otani A, Mineoka A, Watanabe H: Inhibition by Long-term Fermented Miso of Induction of Pulmonary Adenocarcinoma by Diisopropanolnitrosamine in Wistar Rats. *Hiroshima J Med Sci*, 52, 9-13, 2003.
- 74) Shiraki K, Kashiwabara S, Kashimoto N, Une K, Yano R, Otani S, Mineoka A, Hashimoto K, Kageyama N, Nishioka T, Kominami Y, Sasaki A, Shiraishi M, Uesaka T, Katoh O, Watanabe H: Inhibition by long-term fermented miso of induction of pulmonary adenocarcinoma by Diisopropanolnitrosamine (BHP) in wister rats. *Miso Tech Sci*. 51, 391-397, 2003.(in Japanese)
- 75) Koo LC, Ho JH, Matsuki H, Shimizu H, Mori T, Tominaga S: A comparison of the prevalence of respiratory illnesses among nonsmoking mothers and their children in Japan and Hong Kong. *Am Re Respir Dis* 138, 290-295,

- 1988.
- 76) Swanson CA, Mao BL, Li JY, Lubin JH, Yao SX, Wang JZ, Cai SK, Hou Y, Luo QS, Blot WJ : Dietary determinants of lung-cancer risk: results from a case-control study in Yunnan Province, China. *Int J Cancer* 50, 876-880, 1992.
  - 77) Ozawa K., Watanabe Y, Ito Y, Suzuki K, Tamakoshi A, Seki N, Takaaki T, Wakai K, Ando M, Ohno Y, for the JACC Study group: Dietary habits and risk of lung cancer death in a large-scale cohort study (JACC Study) in Japan by sex and smoking habit. *Jpn J Cancer Res* 92, 1259-1269, 2001.
  - 78) Ito A, Watanabe H, Basaran N: Effects of soy products in reducing risk of spontaneous and neutron-induced liver tumors in mice. *Int J Oncol*, 2, 773-776, 1993.
  - 79) Ogundigie PO, Roy G, KaninG, Goto T, Ito A: Effect of biochanin A or testosterone on liver tumors induced by a combined treatment of DEN and fission neutron in BCF1 mice. *Oncol Rep*, 2, 271-275, 1995.
  - 80) Kurozawa H, Ogimoto I, Shibata A, Nose T, Yoshimura T, Suzuki H, Sakata R, Fujita Y, Ichikawa S, Iwai N, Fukuda K, Tamakoshi A: Dietary habits and risk of death due to hepatocellular carcinoma in a large scale cohort study in Japan. Univariate analysis of JACC study data. *Kurume Med J*, 51, 141-149, 2004.
  - 81) Sharp GB, Lagarde F, Mizuno T, Sauvaget C, Fukuhara T, Allen N, Suzuki G, Tokuoka S: Relationship of hepatocellular carcinoma to soya food consumption: A cohort-based case-control study in Japan. *Int J Cancer*, 115, 280-295, 2005.
  - 82) Yamamoto S, Sobue T, Kobayashi M, Sasaki S, Tsugane S: Soy, isoflavones, and breast cancer risk in Japan. *J Natl Cancer Inst*, 95, 906-913, 2003.
  - 83) Baggott JE, Ha T, Vaughn WH, Juliana M, Hardin JM, Grubbs CJ: Effect of Miso (Japanese Soybean paste) and NaCl on DMBA-induced rat mammary tumors. *Nutr Cancer*, 14, 103-109, 1990.
  - 84) Gotoh T, Yamada K, Ito A, Yin H, Kataoka T, Dohi K: Chemoprevention of N-nitroso-N-methylurea rat mammary cancer by miso and tamoxifen, alone and in combination. *Jpn J Cancer Res*, 89, 487-495, 1998.
  - 85) Yamagishi K, Iso H, Tanigawa T, Cui R, Kudo M, Shimamoto T: High sodium intake strengthens the association between angiotensinogen T174M polymorphism and blood pressure levels among lean men and women: a community-based study. *Hypertens Res.*, 27, 53-60, 2004.

- 86) Tamaki J, Kikuchi Y, Yoshita K, Takebayashi T, Chiba N, Tanaka T, Okamura T, Kasagi F, Minai J, Ueshima H; HIPOP-OHP Research Group: Stage of changes for salt intake and urinary salt excretion: Baseline results from the high-risk and population (HIPOP-OHP) study. *Hypertens Res*, 27, 157-166, 2004.
- 87) Tochikubo O, Nishijima K: Sodium intake and cardiac sympatho-vagal balance in young men with high blood pressure. *Hypertens Res*, 27, 393-398, 2004.
- 88) He FJ, MacGregor GA: Blood pressure--importance of salt intake. *Am J Hypertens*, 18, 1258-1259, 2005.
- 89) He FJ, Markandu ND, MacGregor GA: Modest salt reduction lowers blood pressure in isolated systolic hypertension and combined hypertension. *Hypertension*, 46, 66-70, 2005.
- 90) He J, Gu D, Wu X, Chen J, Duan X, Chen J, Whelton PK: Effect of soybean protein on blood pressure: a randomized, controlled trial. *Ann Intern Med*, 143, 1-9, 2005.
- 91) Obarzanek E, Velletri PA, Cutler JA: Dietary protein and blood pressure. *JAMA*, 275, 1598-1603, 1996.
- 92) Watanabe H, Kashimoto N, Kajimura J, Kamiya K: A miso (Japanese soybean paste) diet conferred greater protection against hypertension than a sodium chloride diet in Dahl salt-sensitive rats. *Hypertension Res*, 29, 731-738, 2006.
- 93) Washburn S, Burke GL, Morgan T, Anthony M: Effect of soy protein supplementation on serum lipoproteins, blood pressure, and menopausal symptoms in perimenopausal women. *Menopause*, 6, 7-13, 1999.
- 94) Burke V, Hodgson JM, Beilin LJ, Giangiulioi N, Rogers P, Puddey IB: Dietary protein and soluble fiber reduce ambulatory blood pressure in treated hypertensives. *Hypertension*; 38, 821-826, 2001.
- 95) Hu FB, Stampfer MJ, Manson JE, Rimm E, Colditz GA, Rosner BA, Hennekens CH, Willett WC: Dietary fat intake and the risk of coronary heart disease in women. *N Engl J Med*; 337, 1491-1419, 1997.
- 96) Nevala R, Vaskonen T, Vehniainen J, Korpela R, Vapaatalo H: Soy based diet attenuates the development of hypertension when compared to casein based diet in spontaneously hypertensive rat. *Life Sci*; 66, 115-124,. 2000
- 97) Nakamura M, Aoki N, Yamada T, Kubo N: Feasibility and effect on blood pressure of 6-week trial of low sodium soy sauce and Miso (fermented soybean paste). *Circ J*, 67, 530-534,. 2003
- 98) Kanda A, Hoshiyama Y, Kawaguchi T: Association of lifestyle parameters

with the prevention of hypertension in elderly Japanese men and women: a four-year follow-up of normotensive subjects. *Asia Pac J Public Health*, 11, 77-81, 1999.

- 99) Fujita T: Low salt diet should be reviewed. Miso soup is the best to prevent hypertension by intake of potassium or magnesium. *Miso News Letter* 65-71, 1995.
- 100) Kennedy AR: Prevention of carcinogenesis by protease inhibitors. *Cancer Res.* 54(7 Suppl), 1999s-2005s, 1994.
- 101) Kennedy AR, Davis JG, Carlton W, Ware JH: Effects of dietary antioxidant supplementation on the development of malignant lymphoma and other neoplastic lesions in mice exposed to proton or iron-ion radiation. *Radiat Res*, 169, 615-625, 2008

Table 1 Number of surviving crypts after X-irradiation<sup>6)</sup>

	0 Gy	7 Gy	8 Gy	10 Gy	12 Gy
MF (Control)	123.7±13.1	97.6±10.8	80.8±8.9	48.7±7.1	30.3±5.8
3-4 days fermented miso		112.5±14.0 <sup>**</sup>	97.3±13.7 <sup>**</sup>	53.0±6.4 <sup>*,a</sup>	41.0±6.5 <sup>**a</sup>
120 days fermented miso		108.4±13.9 <sup>**</sup>	84.4±11.7	55.1±5.4 <sup>**a</sup>	43.9±5.1 <sup>**a</sup>
180 days fermented miso	125.2±12.7	103.0±11.6	87.4±9.1 <sup>**</sup>	68.5±9.3 <sup>**</sup>	50.0±5.2 <sup>**</sup>

(Mean±SD)

<sup>\*\*</sup>: Significantly different from MF value (p<0.01)<sup>a</sup>: Significantly different from 180 days fermented miso value (p<0.01)

Table 2 Incidence of gastric tumor by miso and NaCl by *N*-methyl-*N'*-nitro-*N*-nitrosoguanidine (MNNG) in CD(SD) rats<sup>15)</sup>

Group	Total tumor (Number per rat)	Glandular stomach			Small intestine			Other tumor
		Incidence	Size (mm)	Number	Incidence	Size (mm)	Number	
MNNG+10%Miso	17/20(85.0%) a (1.15±0.75) <sup>a</sup>	9/20(45%)	2.3±3.5	0.6±0.7	9/20(45%)	8.2±13.9	0.6±1.0	Sarcoma
MNNG+2.2% NaCl	17/19 (89.5%) <sup>a</sup> (1.32±0.67) <sup>a</sup>	13/19(68%) <sup>a</sup>	4.1±5.1	1.5±0.5	11/19(58%) <sup>a</sup>	11.4±17.1	0.6±0.8 <sup>a</sup>	Squamous cellcarcinoma Plasmacytoma
MNNG+5%Miso	10/19 (52.6%) (0.63±0.68)	7/19(37%)	2.0±2.9	0.5±0.7	2/19(11%) b	5.0±17.5	0.2±0.5	Lymphoma Hepatoma
MNNG+1.1%NaCl	15/20(75.0%) a (0.9±0.64)	12/20(60%)	4.5±4.4 <sup>c</sup>	0.7±0.7	6/20(30%)	4.1±9.1	0.3±0.6	Sarcoma
MNNG+MF	10/19(53.0%)	6/19(32%)	1.2±2.3	0.7±1.1	3/19(16%)	7.6±20.6	0.2±0.4	Plasmacytoma

(0.65±0.67)

ma

Sarcoma

---

(mean±SD)

\*( ) Number of tumor per rat

<sup>a</sup>: Significantly different from MNNG+MF value (P<0.05);

<sup>b</sup>: Significantly different from MNNG+10%*miso* value (P<0.05)

<sup>c</sup>: Significantly different from MNNG+MF value (P<0.01)

Table 3 Inhibition by 180-days fermented miso induction of gastric tumor's number and size by MNNG in CD rats<sup>17)</sup>

Group	Gastric tumor <sup>1</sup>	Adenocarcinoma <sup>2</sup>	
		Number	Size (mm)
MF	5.17±4.45	11	8.0±2.9
3-4 days fermented miso	3.62±6.79	6	11.5±7.6
120 days fermented miso	2.42±2.68	6	4.3±3.6 <sup>a</sup>
180 days fermented miso	1.78±3.34 <sup>*</sup>	9	2.9±4.6 <sup>*,b</sup>
NaCl	3.23±4.08	8	5.1±4.1 <sup>a</sup>

(mean±SD)

<sup>1</sup>. Included negative animal<sup>2</sup>. Positive animal values<sup>\*</sup>: Significantly different from MF value (P<0.05)<sup>a</sup>: Significantly different from early-term value (P<0.05)<sup>b</sup>: Significantly different from early-term value (P<0.01)

Table 4 Effects of dietary miso, NaCl and others on azoxymrthane (AOM)-induced aberrant crypt foci (ACF) in F344 rat colons<sup>33-35)</sup>

Group	ACF/colon	AC/colon	AC/focus
MF (Control)	136.4±41.3	221.4±61.9	1.63±0.10
4.4% Sea salt	39.8±22.0**	62.9±38.1**	1.69±0.48
4.4% Rock salt	57.6±26.8**	98.3±48.1**	1.70±0.17
4.4% Home salt	35.5±11.6**	61.2±21.5**	1.71±0.10
4.4% Coarse salt	74.1±33.5**	123.9±54.1**	1.70±0.14
4.4% Pure NaCl	100.4±62.9	159.8±95.6*	1.61±0.23
2.2% Pure NaCl	115.9±65.2	199.5±92.6	1.82±0.32
20% Miso	66.3±41.1**	124.4±78.0**	1.88±0.18
10% Miso	85.6±70.5*	163.0±130.8	1.94±0.15
5% Miso	134.1±52.2	240.9±92.9	1.83±0.17
3.0% kojic acid	97.5±46.6*	169.9±72.6	1.79±0.20
1.5% kojic acid	91.6±31.3*	167.2±53.4*	1.84±0.20
5ppm Biochanin A	131.0±63.0	226.6±90.0	1.81±0.27

(Mean±SD)

\*: Significantly different from Control value (P<0.05)

\*\* : Significantly different from Control value (P<0.01)

Table 5 Decrease by 180-day fermented miso of colon carcinoma's size induced by AOM in F344 Rats<sup>47)</sup>

Group	Number of tumor	Size (mm)	Size of well-differentiated adenocarcinoma (mm)	Size of total tumor (mm)
2-year fermented miso+AOM	0.50ear f	4.60ear f	9.72±5.43	11.40.6.37
180 days fermented miso+AOM	0.85days	2.72days <sup>a</sup>	3.56±2.56 <sup>b</sup>	3.56days <sup>a,**</sup> 9
3-4 days fermented miso+AOM	0.95days	1.79days	3.66±2.45 <sup>b</sup>	7.97days
MF+AOM	1.10OMys	3.76OMy	6.86±3.38	7.26OMys

(mean±SD)

a: Significantly different from 2 year fermented miso+AOM value (P&lt;0.05)

b: Significantly different from 2 year fermented miso+AOM value (P&lt;0.01)

Table 6 Inhibition by 180-days fermented miso of induction of pulmonary tumor by N-nitrosobis(2-hydroxypropyl)amine (BHP) in Wistar rats<sup>73)</sup>

Group	No of animals	Adenoma	Adenocarcinoma	Tumor with PCNA-strongly positive
BHP	8	62.6A-str	6.56A-s	21.0A-st
BHP+3-4 days fermented miso	15	61.6±19.0	6.3±2.7	15.4±6.6*
BHP+180 days fermented miso	14	57.8±24.9	2.9±1.7**	11.7±4.7**

(mean±SD)

\*\* : Significantly different from BHP value (P<0.01)

Table 7 Blood pressure in male rats<sup>92)</sup>

Systolic blood pressure					
Group	0 week	2 weeks	4 weeks	8 weeks	12 weeks
Dahl Miso(2.3%)		124.2±10.3	144.0±13.0	158.8±20.8	158.8±19.4
Dahl NaCl(2.3%)		131.4±11.4	147.8±9.6	181.0±15.0 <sup>**</sup> , b	191.6±13.4 <sup>**</sup> , b
Dahl NaCl(1.9%)		134.2±10.1	151.6±9.6	181.2±19.2 <sup>**</sup> , b	189.8±30.2 <sup>**</sup> , b
Dahl MF	102.9±11.3	140.1±15.9	145.3±14.2	154.3±7.6	154.6±7.9
SD Miso		117.6±15.8	130.0±8.7	135.5±16.1	132.5±15.0
SD NaCl(2.3%)		127.4±14.2	141.0±17.7	142.8±12.0*	136.6±7.2
SD NaCl(1.9%)		117.0±13.8	144.6±12.5*	132.0±16.6	129.4±6.5
SD MF	95.4±11.5	127.5±7.5	129.3±13.4	130.1±11.8	137.1±18.3
Diastolic blood pressure					
Group	0 week	2 weeks	4 weeks	8 weeks	12 weeks
Dahl Miso(2.3%)		97.7±8.2	112.6±7.6	127.6±21.6	119.9±18.0
Dahl NaCl(2.3%)		107.8±13.5	121.2±6.2 <sup>**</sup> , a	148.4±13.2 <sup>*</sup> , a	166.8±26.9 <sup>**</sup> , b
Dahl NaCl(1.9%)		104.8±18.9	120.2±6.8a	152.8±25.2 <sup>*</sup> , a	165.8±15.1 <sup>**</sup> , b
Dahl MF	77.5±11.6	108.3±9.8	113.8±13.5	127.9±7.8	123.9±8.4
SD Miso		84.6±10.7	101.3±9.5	102.5±9.7	101.0±10.6
SD NaCl(2.3%)		93.2±10.4	104.8±15.7	110.4±19.6	101.8±7.7
SD NaCl(1.9%)		80.8±11.2	112.0±19.2	106.2±12.5	101.2±8.8
SD MF	68.7±8.5	93.6±9.5	97.1±13.8	102.4±7.5	112.2±14.1

\*: Significantly different from same strain MF (P<0.05), \*\*: Significantly different from same strain MF (P<0.01)

<sup>a</sup>: Significantly different from same strain Miso (P<0.05)<sup>b</sup>: Significantly different from same strain Miso (P<0.01)

Table 8 Blood pressure in female rats<sup>92)</sup>

Systolic blood pressure						
Group	0 week	2 weeks	4 weeks	8 weeks	12 weeks	
Dahl Miso		121.9±10.8	134.4±11.4	145.0±12.9	143.1±16.0	
Dahl NaCl(2.3%)		118.4±8.4	153.0±5.6** b	161.4±7.6** b	173.6±15.8** b	
Dahl NaCl(1.9%)		114.0±10.0	141.4±14.5	142.8±19.2	146.6±8.0	
Dahl MF	108.3±12.2	119.5±10.0	137.4±12.2	140.6±10.5	143.8±11.6	
SD Miso		114.9±11.7	133.5±7.0	131.7±12.9	132.2±12.1	
SD NaCl(2.3%)		116.0±14.4	139.4±9.1*	136.8±10.9*	142.4±7.9 <sup>a</sup>	
SD NaCl(1.9%)		115.8±12.4	130.4±3.0	126.8±15.3	133.8±8.8	
SD MF	95.9±11.5	115.7±8.6	124.8±12.2	124.2±10.9	132.0±16.8	
Diastolic blood pressure						
Group	0 week	2 weeks	4 weeks	8 weeks	12 weeks	
Dahl Miso		93.8±5.7	107.7±12.7	112.3±10.5	110.9±23.1	
Dahl NaCl(2.3%)		97.2±10.4	116.0±13.8	123.0±8.8** b	141.8±7.3** b	
Dahl NaCl(1.9%)		82.8±21.2	116.2±12.2	111.4±11.9	120.4±6.9	
Dahl MF	80.0±13.9	92.8±12.9	104.7±14.5	109.1±11.8	109.9±18.8	
SD Miso		85.9±8.5	108.4±7.5	99.2±12.6	99.4±5.5	
SD NaCl(2.3%)		88.2±4.8	106.6±11.1	114.4±11.3** b	111.6±13.0	
SD NaCl(1.9%)		86.6±13.6	101.4±6.1	95.0±10.4	104.2±14.2	
SD MF	67.2±10.3	87.5±10.8	95.6±9.9	98.7±12.4	103.4±13.7	

\*:Significantly different from same strain MF (P<0.05); \*\*:Significantly different from same strain MF (P<0.01)

<sup>a</sup>: Significantly different from same strain Miso (P<0.05); <sup>b</sup>: Significantly different from same strain Miso (P<0.01)

